

## THE HYPOTHETICAL SECONDARY BRAIN OF THE DIGESTIVE APPARATUS

JANOS VINCZE, GABRIELLA VINCZE-TISZAY

Health Human International Environment Foundation,  
Budapest, Hungary

### ABSTRACT

Digestion is the complex physiological process by which food that has entered the alimentary tract undergoes physical and chemical changes and the nutrients in it are absorbed into the blood and lymph. Starting from Bertalanffy's systems theory, we wrote the differential equation systems of the digestive tract. We need to consider the digestive system as a so-called hierarchical system, because it has many subsystems. The digestive system has a two-level control. Block diagrams can be used to examine the kind of goal orientation we want to make effective because this always depends on the current needs of the given organism. A block diagram of the brain-controlled digestive tract in case of normal functioning was recorded. A block diagram of the part directed by the hypothetical secondary brain was also recorded. We note that this part is functioning also under normal circumstances, only the brain activity obscures (masks) the everyday activity of the hypothetical brain.

**Keywords:** digestive apparatus, systems theory, block diagram, biophysical modeling, and hypothetical secondary brain

### 1.0 INTRODUCTION


Digestion has entered the alimentary tract and undergoes physical and chemical changes. The physical changes in food consist in its mechanical processing by which it is grounded, mixed, and dissolved. The chemical changes comprise a number of successive stages of protein, fat, and carbohydrate hydrolysis, occurring under the influence of hydrolytic enzymes. [1] Three groups of these enzymes are distinguished: 1) those responsible for the breakdown of proteins, or proteases; 2) those responsible for the breakdown of fats, or lipases; and 3) those responsible for the breakdown of carbohydrates, or carbohydrases. The enzymes are produced in the secretory cells of the glands concerned with digestion and enter the alimentary canal as constituents of saliva and of the gastric, pancreatic, and intestinal juices. Any one nutrient is attacked by the various groups of enzymes, one acting after the other, and is gradually broken down into less complex chemical compounds.

Most of the substances contained in food—proteins, fats and carbohydrates are high-molecular compounds that cannot be absorbed and utilized by the cells of the organism without being chemically treated in the alimentary tract. [2] Only the simplest compounds, readily soluble

in water and devoid of species specificity, which are formed from them, can penetrate the wall of the alimentary tract into the blood. All the functions of the alimentary organs are regulated by complex nervous and humoral mechanisms.

## 2.0 SYMBIOSIS OF THE STRUCTURE AND FUNCTION

A person consumes many times their body weight in the form of food during the decades of their life. Meanwhile, from infancy, their body grows 15 to 20 times. [3] According to the mechanical preparation, delivery, chemical transformation and circulation of food, the basic functions of the digestive system can be grouped around movement activity, fluid secretion and digestion, as well as absorption, which are not independent functional groups but take place in a closely and mutually coordinated manner. (Fig. 1.)



**Figure 1. The process-diagram – after the requirement of structure-function – in the human digestive apparatus:**

The digestion process should be optimized, we are not eating continuously, but intermittently, approx. 3–4 times a day, we do not eat unlimited quantities, because the body does not require it, etc. Defecation should be made possible 1–2 times a day. So optimization means that fluctuations around the minimum and maximum values should be characteristic of the daily digestive process. There is a whole other kind of external restriction, namely the amount of money we have to buy certain foods, and the kinds of foods we have in our environment. Another external barrier may be whether we can dedicate enough time for meals in our daily activities. [3]

The more frequent diseases of the digestive apparatus are: gastro-duodenal ulcer, gastric ulcer, gastric cancer, hepatitis, liver cirrhosis, hepatic, gallbladder, colon cancer, chronic colitis, Crohn's disease, irritable colon – colopathy, chronic gastritis, digestive haemorrhage. [5, 6]The disturbances are seldom accentuated by stress.

Hyperponderability is not related only to the digestive system. [7] Women tend to gain weight during menopause – an average of 5–10 kg, with the modification of fat spread, which deposits mainly on the abdomen and hips. [8] This weight gain can be partially explained by hormonal factors [9] (the centres of appetite and satiety are in the hypothalamus, near the ones adjusting the ovarian activity); it is favoured also by the decrease of physical activity related to age, to stress [10] – personal or professional – which lead to an excessive feeding, to searching in food – and especially sweets – a compensation for frustrations.

### **3.0 THE DIGESTIVE APPARATUS IS SUBSYSTEM OF ORGANISM**

Starting from Bertalanffy's systems theory, we wrote the differential equation systems of the digestive tract. We need to consider the digestive system as a so-called hierarchical system, because it has many subsystems. Let us examine the digestive tract from the point of view of systems theory. We have to assume that the digestive tract as a system is an integrated whole, even though it derives from various structures and subfunctions. [11] The other starting point should be that as a system it has a certain objective, and that the balance of these can vary greatly. The processes taking place in the digestive tract aim to optimize the functions of the system as a whole according to the objective, and to achieve maximum compatibility of the digestive subsystems. The main goal is to make the whole system work, not for a subset to operate optimally. [12] A given system (digestive tract), with its core inputs and outputs, is a subset of a larger "super system", the human organism. The requirements for different levels of subsystems are not the same, for example, think of two subsystems: the activity of the parotid gland and the pancreas. Subfunctions can enable the overall operation, and they contribute to the success of the entire digestion. Thus, the subfunctions facilitate the functioning of the digestive tract as a whole. [13]

So, for example, the kind of food we take in affects the attitude of the subsystems and their sub-functions to this food. For example, they have a qualitatively different attitude towards liquid foods than towards solid foods, but they also have a different attitude when consuming sugary or fibrous foods. The lifetime of digestion depends on the quality of the food, the intensity of the small and large intestine sub-functions, etc. [14]

People actually plan their meals, and buy their daily food accordingly. If we look at all this at system level, it can be illustrated in the form of a block diagram (Fig.1.):

The digestive system has a two-level control. Block diagrams can be used to examine the kind of goal orientation we want to make effective because this always depends on the current needs of the given organism. A block diagram of the brain-controlled digestive tract in case of normal functioning was recorded.

### **4.0 THE TWO-LEVEL BIOPHYSICAL CONTROL OF THE DIGESTIVE TRACT IN HUMAN ORGANISM**

The central nervous system regulation of the digestive tract is primarily achieved through the nerves in the brain. There is no center in the brain that controls digestion, and that would control all processes of the digestive tract from a single center. [15]

Consider the cerebral nerves of the 12 pairs that work at different levels in the digestive tract:

V. The trigeminal nerve is mixed, with three branches: the ophthalmic, the maxillary and the mandibular branch. The first two are sensory, the third one is mixed. It innervates the facial skin and masseter muscles.

VII. The facial nerve is mixed, providing the innervation of the mimic muscles, the sensitivity of taste, the function of the salivary glands from under the tongue and the jaw, and the secretory activity of the tear glands.


IX. The glossopharyngeal nerve is mixed, providing the innervation of laryngeal muscle, taste perception and the salivation of the parotid.

X. The vagus nerve contains mixed, sensory, motor and vegetative fibers: it controls most of the internal organs.

XII. The nerve under the tongue is a motor that affects the tongue muscles.

The digestion process is described in terms of equations, symbols, circuit diagrams, as a result of which requirements can be concretely implemented in an acceptable solution. (Fig. 2.)

On the basis of systems theory, every subsystem of a system has a certain level of integrity, with its own structural and functional stability. In our opinion, this is true of the human body's digestive apparatus as the subsystems of the organism as well. [16, 17]



**Fig. 2. Process diagram of the digestive tract according to the requirements of function and structure**

The concept of the subsystem purports that it must have control over its own structure. It follows, in our opinion, that digestive apparatus of the human organism contain a so-called “hypothetical secondary brain”, which is responsible for certain control functions over the

particular apparatus. (Fig. 3.) [15] The hypothetical secondary brain of the digestive apparatus is probably located in the large intestine.




Fig. 3. The block diagram of the digestive tract – only „hypothetical seconder brain” – control

**CONCLUSIONS**

The digestive apparatus is made of organs in which the digestion of the foods and their absorption takes place. At the same time, in the last segment of the digestive tube, the rectum, the elimination of the unabsorbed rests takes place through the act of defecation. [18] Eating

is strongly influenced by the unique past and mental state of the organism. It has never consumed certain foods (which did not exist in its environment, or did not like them); or a previous consumption of these led to illness or allergic condition.

The nutrition diseases are infections caused by disturbances in the metabolism of the food substances. [19] The metabolic deviations have multiple causes: exogenous, endogenous and mixed. The most important exogenous factor is the food intake. From the exogenous factors the most important are: the constitutional, hereditary and endocrine dysfunctions.

We wrote the differential equation systems of the digestive tract by system theory and characterized its activity by block diagram.

In our opinion, this “hypothetical secondary brains” of the digestive apparatus function continuously over a person’s life, but its control function is so weak that due to the dominant activity of the central nervous system, it has not yet been detected and is still uncovered by scientific research. [15]

A fortunate model describes the given momentary acquisitions but beyond this it enables assumptions which can be then translated into the language of the experiment and on this basis you can arrange certain new experimental conditions and situations. An adequately selected model will promote not only raising the question but also the scientific investigation of the phenomena. Forming a biophysical model [19] is not the task of the biologist or physicist alone, a good model can successfully be constructed only by common, collective-work. This is typically the task of the biophysicist and a problem which falls within the competence of this discipline. Model is always an approximation, the user of the model has to take into consideration that he can approach only the absolute truth just through the endless series of relative truths.

## REFERENCES

- Lin, J. T.: Screening of gastric cancer, who, when, and how. *Clin. Gastroenterol. Hepatol.*, 2014; 12: 135–138.
- Bernal, W., Wendon, J.: Acute liver failure. *N. Engl. J. Med.*, 2013; 369: 2525-2534.
- Lazaridis K. N., LaRusso, N. F.: Primary sclerosing cholangitis. *Lancet*, 2015; 386: 1565-1568.
- Louvet A., Mathurin, P.: Alcoholic liver disease: mechanisms of injury and targeted treatment. *Nat. Rev. Gastroenterol. Hepatol.*, 2015; 12: 231-240.
- Hart, P. A., Zen, Y., Chari, S. T.: Recent advances in autoimmune pancreatitis. *Gastroenterology*, 2015; 149: 39–51.
- Graham, C., Mullen A., Whelan K.: Obesity and gastrointestinal microbiota: a review of associations and mechanisms. *Nutr. Rev.* 2016; 73: 376-382.

Roberts, D. L., Dive C., Renehan A. C.: Biological mechanisms linking obesity and cancer risk: new perspectives. *Annu. Rev. Med.* 2010; 61: 301-307.

Wadden, T. A., Sternberg, J. A., Letizia, K. A.: Treatment of obesity by very low calorie diet, behavior therapy, and their combination: a five-year perspective. *Int. JOb.*, 1989; 13: 39-45.

Casals-Casals, C., Desvergene B.: Endocrine disruptors: from endocrine to metabolic disruption. *Annu. Rev. Phys.* 2011; 73: 135-141.

Starcke, K., Brand, M.: Effects of Stress on Decision Under Uncertainty. *Psychological Bull.* 2016; 1323: 151–181.

Vincze, J. (2015): *Biophysics of Digestive System*. NDP P., Budapest

Vincze, J. (2007): *Interdisciplinarity*, NDP P., Budapest

Vincze, J. (2018). *Medical Biophysics*. NDP P., Budapest

Vincze, J.: The Biophysical Modeling of the Evaluation of the Laboratorial Diagnosis Zones. *American Journal of Internal Medicine.* 2020; 8(1): 1–7.

Vincze, J.: The “hypothetical secondary brain”, *Medical Research Archives.* 2019; 7(7): 1–3.

Vincze J.: (2020) *The Biophysics of the Apparatuses in the Human Organism*. NDP P., Budapest,

Vincze, J.: (2020) *The Biophysical Modeling of the Apparatuses in the Human Organism*. Lambert, Berlin.

Paumgatner G. M.: Biliary physiology and disease reflections of a physician-scientist. *Hepatology*, 2010; 51:1095-1110.

Vincze, J. (2021). *Biophysical vademecum*. NDP P., Budapest